BaylorScott&White
ORTHOPEDIC ASSOCIATES OF DALLAS
A member of HealthTexas Provider Network

Manoj P. Reddy, MD
Sports Medicine & Shoulder Surgery
Orthopedic Associates of Dallas
469.800.7200 office | 469.800.7210 fax
DallasOrtho.com | ReddyOrtho.com

Arthroscopic Rotator Cuff Repair

REHABILITATION PROTOCOL

PHASE	RANGE OF MOTION	IMMOBILIZER	THERAPEUTIC EXERCISE
Phase I **0–6 weeks** GOAL TO ADVANCE *Repair protected; pain controlled*	NO shoulder motion, passive or active. No reaching overhead or behind back	Sling with abduction pillow at all times, including sleep, except hygiene	Hand/wrist/elbow active ROM, scapular mobility in sling, ice
Phase II **6–10 weeks** GOAL TO ADVANCE *Passive flexion 100-120°, ER 25-45°*	Passive with therapist: supine elevation 0-100°, seated ER 0-30°, table slides, pendulums. No active motion; no reaching behind back	Wean with physician clearance	Scapular retraction, elevation and depression out of sling. No weight bearing through the arm
Phase III **10–18 weeks** GOAL TO ADVANCE *Active flexion >120° without shrug*	Week 10: active-assisted (cane) lying down → reclined (week 11) → upright (week 12). Active standing ER week 12; active elevation week 14	None	Weeks 14-18: gentle (submaximal) isometrics with arm at side, standing rows. Grade I-II joint mobilizations after week 10
Phase IV **18–22 weeks** GOAL TO ADVANCE *Full motion; pain-free strengthening*	Full motion expected. Stretching: pec, internal rotation with towel, cross-body, sleeper	None	Prone I/Y/T/W, band ER/IR, rows, punches, biceps/triceps. Lift ≤5 lb. No straight-arm side raises or "empty can"
Phase V **22–26 weeks** GOAL TO ADVANCE *4+/5 pain-free strength*	Full; more aggressive stretching if needed	None	ER/IR at 45° and 90° abduction, full can (1-2 lb), diagonal patterns, push-up progression. Lift ≤10 lb; no overhead lifting
Phase VI **26–30 weeks** GOAL TO ADVANCE *85-90% strength vs other side*	Full without discomfort	None	Gym program, work- and sport-specific training**

NOTES

**If approved by physician

This protocol is a guideline. Your surgeon may adjust it based on your surgery and progress.
Call the office for fever, wound drainage, calf pain or swelling, or new numbness or tingling.

Questions: 469.800.7200
Rev. 10/2026